

STIC-ILL

NO 409008

From: Steadman, David (AU1652)  
Sent: Tuesday, August 20, 2002 1:10 PM  
T: STIC-ILL  
Subject: literature request for 09/815,533

Art Unit: 1652  
Office: 10D-04  
Mailbox: 10C-01 M3  
Case Serial #:09/815,533

Please provide the following references:

7870493

- 1) J Chromatogr 1990 Feb 23;525(2):297-306  
Purification of urokinase by combined cation exchanger and affinity chromatographic cartridges.  
Hou KC, Zaniwski R.
- 2) Purification of high-molecular-weight and low-molecular-weight urokinase and kinetic study  
Sun, Tian-Xiao; Wang, Hong-Mei; Xu, Chang-Fa  
Shengwu Huaxue Zazhi (1997), 13(3), 344-349
- 3) Isolation, purification and comparative studies of certain properties of high- and low-molecular-weight urokinases of human urine.  
Sun, Leqin; Zhang, Hongzu; Zhu, Dexu  
Shengwu Huaxue Yu Shengwu Wuli Xuebao (1984), 16(3), 303-6
- 4) J Biochem (Tokyo) 1981 Jul;90(1):225-32  
A comparative study of high molecular weight urokinase and low molecular weight urokinase.  
Nobuhara M, Sakamaki M, Ohnishi H, Suzuki Y.
- 5) Enzyme 1981;26(4):221-4  
Kinetic studies of three different molecular forms of urokinase for the activation of native human plasminogen.  
Toki N, Takasugi S, Sumi H.
- 6) Thromb Haemost 1983 Apr 28;49(2):91-5  
Purification of high molecular weight urokinase from human urine and comparative study of two active forms of urokinase.  
Shibatani T, Kakimoto T, Chibata I.
- 7) Thromb Haemost 1982 Jun 28;47(3):197-202  
Rapid isolation of high molecular weight urokinase from native human urine.  
Huber K, Kirchheimer J, Binder BR.
- 8) Chem Pharm Bull (Tokyo) 1981 Feb;29(2):463-71  
Comparative studies on two active enzyme forms of human urinary urokinase. I. Purification by serial column chromatography and homogeneity analyses of molecular weight and isoelectric point.  
Miwa N, Takayanagi H, Suzuki A.

6788545

Thank you,  
David J. Steadman  
Art Unit 1652  
CM1, 10D-04  
308-3934

son, S.; Westman, S.: Mast cell con- acid metabolism in the epididymal ese mice. *Acta physiol. scand.* 58: 3).

1, E.: Methods for determination of e in man and animals. *J. Lipid. Res.* 9: 68).

cerolipid biosynthesis in rat adipose n. *J.* 170: 153-160 (1978).

piro, D.; Fallon, H.J.: Triacylglyc- is in adipose tissue of the obese- c mouse. *Biochem. J.* 158: 327-334

coff, M.A.; Bell, R.M.: Triacylglyc- in isolated fat cells. An effect of insu- mal fatty coenzyme A ligase activi- m. 251: 1488-1492 (1976).

ty acid synthesis in adipose tissue itiated water. *Biochemistry* 7: 3708-

ction of insulin effector system of cells by proteolytic enzymes. *J. biol.* 177-5784 (1969).

es, A.B.: Metabolism of isolated fat lar inhibitory actions of phospholi- dium perfringens  $\alpha$  toxin and lipo- s and theophylline. *J. biol. Chem.* (1966).

owski, M.J.; Segal, R.: Effect of insu- llular character of rat adipose tissue. 13: 616-623 (1972).

erti, K.G.G.M.; Williamson, D.H.: tozotocin on carbohydrate and lipid the rat. *Endocrinology* 89: 827-834

aggerson, E.D.: Interactions of insu- aline with glycerolphosphate acyla- in fat-cells from rat. *FEBS Lett.* 64:

herjee, C.; Jungas, R.L.: Studies on n of activation of adipose tissue py- ogenase by insulin. *J. biol. Chem.* 247: 373).

tember 30, 1980  
ember 5, 1980

mdar, Medical Research Institute,  
ite of Technology, 3325 W. New  
e, Melbourne, FL 32901 (USA)

*Enzyme* 26: 221-224 (1981)

## Kinetic Studies of Three Different Molecular Forms of Urokinase for the Activation of Native Human Plasminogen

Naotika Toki<sup>a</sup>, Sumiyoshi Takasugi<sup>b</sup>, Hiroyuki Sumi<sup>c</sup>

<sup>a</sup>Department of Dermatology, and <sup>b</sup>Department of Surgery, Hiroshima University, School of Medicine, Japan; <sup>c</sup>Michael Reese Research Foundation, Chicago, Ill., USA

**Key Words.** Urokinase · Glu-PG · Lys-PG · Michaelis constant · Catalytic constant

**Abstract.** The kinetic parameters of three different molecular forms of urokinase (UK) for the activation of native Glu-plasminogen were compared. The apparent Michaelis constant ( $K_m$ , app.) of each UK was almost of the same order of magnitude ( $31-38 \mu M$ ), but the catalytic constants ( $k_c$ ) were observed to be different: UK<sub>h</sub> (high molecular weight form, molecular weight 53,000),  $2.4 \pm 0.2 s^{-1}$ ; UK<sub>l</sub> (low molecular weight form, molecular weight 33,000),  $0.83 \pm 0.10 s^{-1}$ , and UK<sub>t</sub> (trypsin-digested form, molecular weight 36,000),  $0.91 \pm 0.18 s^{-1}$ . The overall second order rate constant,  $k_c/K_m$  calculated for UK<sub>h</sub> was  $7.7 \times 10^4 M^{-1} s^{-1}$ , higher than for UK<sub>l</sub> ( $2.2 \times 10^4 M^{-1} s^{-1}$ ) or UK<sub>t</sub> ( $2.4 \times 10^4 M^{-1} s^{-1}$ ), indicating the possibility of a much higher degree of enzymatic specificity and efficiency.

### Introduction

Christensen and Müllertz [1] and Christensen [2] compared the kinetics of urokinase (UK) in the activation of native Glu-plasminogen (Glu-PG) and its partially degraded form (Lys-PG), and concluded that the binding characteristics (the Michaelis constant,  $K_m$ ) are identical for the two molecules, but their catalytic rates ( $k_c$ ) are significantly different (Lys-PG  $\gg$  Glu-PG). Recently, a few authors have demonstrated the difference in thrombolytic effect and PG activation activity by different UK molecules using the fibrin plate method [3, 4], Chandler's loop method

[3], and sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis [5].

In this report, the kinetic parameters of three different molecular forms of UK for the activation of native Glu-PG were compared in purified systems.

### Materials and Methods

All materials used were of reagent grade quality. The determination methods of protein and UK activity, and immunoelectrophoresis were the same as described previously [6, 7]. The molecular weight of UK was determined using SDS polyacrylamide gel electrophoresis according to the method of Weber and Osborn [8].




Fig. 1. SDS polyacrylamide gel electrophoresis of purified UK. Sample diluent consisted of 6.0 mol/l urea, 1% SDS and 0.01 mol/l sodium phosphate buffer at pH 7.0. a = Reference proteins of bovine serum albumin, egg albumin, chymotrypsinogen A and cytochrome c; b = UK<sub>h</sub>; c = UK<sub>i</sub>; d = UK<sub>t</sub>, respectively. The numbers on the right indicate the molecular weight in thousands.

**Urokinase.** Three different molecular forms were prepared: (1) UK<sub>h</sub>, molecular weight 53,000, which is assumed to be the native UK form [6, 9–13], was purified from a commercial preparation (The Green Cross Co.) by affinity chromatography on [N<sup>α</sup>-(ε-aminocaproyl)-DL-homoarginine hexylester]-Sephadex column followed by Sephadex G-100 gel filtration, as described previously [6]. (2) UK<sub>i</sub>, molecular weight 33,000, which is the modified UK form, was also purified by the same method as UK<sub>h</sub> from a commercial preparation (Mochida Pharmaceutical Co.) as described previously [6]. (3) UK<sub>t</sub>, molecular weight 36,000, which is a trypsin-digested form, was prepared from UK<sub>h</sub> by a method based on that of Lesuk et al. [11]. 0.2 ml of trypsin (Sigma Chemical Co.; 0.86 μg/ml of 0.05 mol/l Tris-HCl, 0.01 mol/l CaCl<sub>2</sub>, pH 8.0) was added to 1.0 ml of UK<sub>h</sub> (2.0 mg/ml of 0.05 mol/l Tris-HCl, 0.1 mol/l NaCl, 0.01 mol/l CaCl<sub>2</sub>, pH 8.0). After the incubation of this mixture for 5 h at room temperature, UK<sub>t</sub> was separated by Sephadex G-100 (Superfine) gel filtration (column: 1.0 × 160 cm). Elution was performed with 0.15 mol/l

ammonium bicarbonate, 0.2 mol/l NaCl, and with that portion which corresponded to the active peak of the partition coefficient  $K_{av}$  [14] 0.39 was collected. Purified preparations were concentrated by ultrafiltration (Collodion-bags, SM 13,200) in the presence of mannitol (1.0%) and lyophilized. They were observed to be homogeneous when subjected to immunoelectrophoresis [6]. In SDS polyacrylamide gel electrophoresis, UK<sub>i</sub> and UK<sub>t</sub> showed a single protein band, but UK<sub>h</sub> showed one main and one minor band (m.w. approx. 100,000, possibly the dimer form of UK<sub>h</sub> or of another contaminating protein) (fig. 1). The specific activities determined by the fibrin plate method [6] were: UK<sub>h</sub> 87,000; UK<sub>i</sub> 79,900, and UK<sub>t</sub> 101,000 international units (IU)/mg of protein, respectively.

**Human Glu-PG.** Purification and determination methods were the same as described previously [1, 2] (>96% Lys-PG as NH<sub>2</sub>-terminal amino acid determined qualitatively by the method of Gros and Labouesse [15]).

**Kinetics.** The kinetic parameters were determined as described by Christensen and Müllertz [1] and Christensen [2]. In a solution containing UK, Glu-PG and α-N-benzoyl-L-arginine ethylester (Bz-Arg-OEt) (Sigma Chemical Co.), Glu-PG is converted to plasmin, which subsequently hydrolyzes Bz-Arg-OEt. The amount of Bz-Arg-OH produced is recorded as a function of time and analyzed to give the corresponding rate of plasmin formation [1]. According to our data (unpublished), 1 mol of UK<sub>h</sub>, UK<sub>i</sub>, and UK<sub>t</sub> contains approximately  $5.5 \times 10^{12}$ ,  $7.3 \times 10^{12}$ , and  $7.0 \times 10^{12}$  IU, respectively. In all kinetic experiments the buffer used was 0.05 mol/l Tris-HCl, 0.1 mol/l NaCl, pH 7.8.

## Results and Discussion

In table I, the calculated kinetic parameters of highly purified UK preparation are shown. The apparent dissociation constants,  $K_m$  of UK<sub>h</sub> and UK<sub>i</sub> were similar ( $31 \pm 8$  and  $38 \pm 6 \mu M$ ) while the catalytic rate constant,  $k_c$ , of UK<sub>h</sub> was approximately 3 times greater than that of UK<sub>i</sub>. All  $K_m$  values are comparable to, but the  $k_c$  values of UK<sub>h</sub> are

Three Diff

Table

Enzyme

UK<sub>h</sub>, molUK<sub>i</sub>, moleUK<sub>t</sub>, mole

Values

consider

[2] ( $K_m$ 

This ma

ecules u

ration (

molecu

generally

weight

and UK

33,000

UK<sub>h</sub> [9,

shown t

molecu

20,000

bonds [

residue

'heavy

UK<sub>i</sub> has

tide cha

phoresis

that hav

the mol

during

comple

[11] fir

dation

lated th

studies

As c

parame

bonate, 0.2 mol/l NaCl, and with 1 corresponded to the active peak of efficient  $K_{av}$  [14] 0.39 was collected. Samples were concentrated by ultrafiltration (SM 13,200) in the presence of sodium chloride and lyophilized. They were observed when subjected to immunoelectrophoresis on SDS polyacrylamide gel electrophoresis. They showed a single protein band, but with two bands (m.w. 33,000 and 53,000, possibly the dimer form of  $UK_h$  or of activating protein) (fig. 1). The specific activity was determined by the fibrin plate method [6]:  $UK_h$  79,900, and  $UK_i$  101,000 (IU)/mg of protein, respectively.  $UK_g$ . Purification and determination were the same as described previously [1, 2] as  $NH_2$ -terminal amino acid determined by the method of Gros and La-

kinetic parameters were determined by Christensen and Müllertz [1] and in a solution containing UK, Glu-PG (L-arginine ethylester (Bz-Arg-OEt) Co.). Glu-PG is converted to plasminogen which hydrolyzes Bz-Arg-OEt. The  $-OH$  produced is recorded as a function of time to give the corresponding formation [1]. According to our data the mol of  $UK_h$ ,  $UK_i$ , and  $UK_g$  contains  $1 \times 10^{12}$ ,  $7.3 \times 10^{12}$ , and  $7.0 \times 10^{12}$  in all kinetic experiments the buffer mol/l Tris-HCl, 0.1 mol/l NaCl,

## Discussion

The calculated kinetic parameters of purified UK preparation are similar to parent dissociation constants, and  $UK_i$  were similar ( $31 \pm 8$ ) while the catalytic rate constant  $k_c$  was approximately 3 times that of  $UK_i$ . All  $K_m$  values are similar but the  $k_c$  values of  $UK_h$  are

Table I. Kinetic parameters of three different UKs for Glu-PG activation

Enzyme	Catalytic constant ( $k_c$ ) $s^{-1}$	Apparent Michaelis constant ( $K_m$ , app.) $\mu M$	Overall rate constant ( $k_c/K_m$ , app.) $M^{-1} s^{-1}$
$UK_h$ , molecular weight 53,000	$k_h = 2.40 \pm 0.2$	$K_h = 31 \pm 8$	$7.7 \times 10^4$
$UK_i$ , molecular weight 33,000	$k_i = 0.83 \pm 0.10$	$K_i = 38 \pm 6$	$2.2 \times 10^4$
$UK_g$ , molecular weight 36,000	$k_g = 0.91 \pm 0.18$	$K_g = 38 \pm 4$	$2.4 \times 10^4$

Values were obtained from four determinations for each example.

considerably higher than those of Christensen [2] ( $K_m = 32 \pm 11 \mu M$ ,  $k_c = 0.26 \pm 0.07 s^{-1}$ ). This may be due to the difference in UK molecules used; Christensen used a crude preparation (10,000 Plouge units/mg) and the molecular variation was not defined. It is now generally accepted that  $UK_h$  with a molecular weight of about 53,000 is the native form, and  $UK_i$  with a molecular weight of about 33,000 is the enzymatically degraded form of  $UK_h$  [9, 10, 12, 13]. Recently,  $UK_h$  has been shown to have two polypeptide chains of a molecular weight of 33,000 (heavy chain) and 20,000 (light chain), linked by disulfide bonds [12, 16]. The active histidine serine residue has been found to be located in the 'heavy chain' [12, 17]. On the other hand,  $UK_i$  has been proven to be a single polypeptide chain by SDS polyacrylamide gel electrophoresis [12, 16]. Despite the many models that have been proposed, the mechanism of the molecular change of  $UK_h$  to  $UK_i$  either during purification or in vivo, is still not completely understood. In 1967, Lesuk et al. [11] first demonstrated the enzymatic degradation of  $UK_h$  in purified systems, and isolated the active molecule,  $UK_i$ . However, no studies of  $UK_i$  have been reported since.

As can be seen in table I, all the kinetic parameters of  $UK_i$  were very similar to those

of  $UK_h$ . From the overall second order rate constant,  $k_c/K_m$  (the fourth column), it may be deduced that the enzymatic efficiency of  $UK_h$  in native PG activation is more than 3 times greater than that in the degraded forms. From the present results and the molecular structures of UK, it can be speculated that the 'light chain' of  $UK_h$  plays a very important role in the activation step of Glu-PG to plasmin.

In recent years, several UK preparations have not only been widely used in the therapy of thromboembolic diseases, but also for enhancement of carcinostatic treatment of malignant tumors. Native UK may prove to be more effective than degraded UKs in such therapy.

## References

- 1 Christensen, U.; Müllertz, S.: Kinetic studies of urokinase-catalyzed conversion of  $NH_2$ -terminal lysine, plasminogen to plasmin. *Biochim. biophys. Acta* 480: 275-281 (1977).
- 2 Christensen, U.: Kinetic studies of the urokinase-catalyzed conversion of  $NH_2$ -terminal glutamic acid plasminogen to plasmin. *Biochim. biophys. Acta* 481: 638-647 (1977).
- 3 Kitamura, M.; Chikamori, K.: Thrombolytic effect of commercial urokinase (Japanese). *Yakkyoku* 25: 1581-1584 (1974).

- 4 Samama, M.; Cazenave, B.; Otero, A.M.: Urokinase I and II activity. *Thromb. Haemostasis* 40: 578-580 (1979).
- 5 Lormeau, J.C.; Coulay, J.; Vairel, E.G.; Choay, J.: A comment on the activities of high and low molecular weight urokinase: fibrinolysis: current fundamental and clinical concepts, pp. 77-82 (Academic Press, New York 1978).
- 6 Sumi, H.; Sasaki, K.; Muramatsu, M.: A simple and rapid purification method of urokinase using a [N<sup>α</sup>-(ε-aminocaproyl)-DL-homoarginine hexylester]-Sepharose column. *Acta haemat. jap.* 41: 766-770 (1978).
- 7 Sumi, H.; Takada, Y.; Takada, A.: The effects of metal ions on esterase activities of urokinase. *Thromb. Diath. haemorrh.* 39: 46-52 (1978).
- 8 Weber, I.; Osborn, M.: The reliability of molecular weight determinations by dodecylsulfate-polyacrylamide gel electrophoresis. *J. biol. Chem.* 244: 4406-4412 (1969).
- 9 Lesuk, A.; Terminiello, L.; Traver, J.H.: Crystalline human urokinase: some properties. *Science* 174: 880-882 (1965).
- 10 White, W.F.; Barlow, G.H.; Mozen, M.M.: The isolation and characterization of plasminogen activators (urokinase) from human urine. *Biochemistry, N.Y.* 5: 2160-2169 (1966).
- 11 Lesuk, A.; Terminiello, L.; Traver, J.H.; Groff, J.L.: Biochemical and biophysical studies of human urokinase. *Thromb. Diath. haemorrh.* 18: 293-294 (1967).
- 12 Johnson, A.J.; Soberano, M.; Ong, E.B.; Levy, M.; Schoellmann, G.: Urinary urokinase. Two molecules or one? *Thrombosis and urokinase*, pp. 59-67 (Academic Press, New York 1977).
- 13 Alkjaersig, N.; Fletcher, A.: Metabolism of urokinase. *Thrombosis and urokinase*, pp. 129-141 (Academic Press, New York 1977).
- 14 Andrews, P.: The gel-filtration behaviour of proteins related to their molecular weights over a wide range. *Biochem. J.* 96: 595-606 (1965).
- 15 Gros, C.; Labouesse, P.: Study of the dansylation reaction of amino acids, peptides, and proteins. *Eur. J. Biochem.* 7: 463-470 (1969).
- 16 Holmberg, L.; Bladh, B.; Åstedt, B.: Purification of urokinase by affinity chromatography. *Biochim. biophys. Acta* 445: 215-222 (1976).
- 17 Ong, E.B.; Johnson, A.J.; Schoellmann, G.: Identification of an active site histidine in urokinase. *Biochim. biophys. Acta* 429: 252-257 (1976).

Received: January 17, 1980

Accepted: November 9, 1980

Naotika Toki, Department of Dermatology,  
Hiroshima, University School of Medicine,  
Kasumi 1-2-3, Hiroshima 734 (Japan)